

A MAYER-VIETORIS THEOREM FOR THE KAUFFMAN BRACKET SKEIN MODULE

WALTER F. LOFARO
*Department of Mathematics,
 The University of Iowa,
 Iowa City, IA 52242, USA
 email: lofaro@math.uiowa.edu*

ABSTRACT

The n th relative Kauffman bracket skein modules are defined and two theorems are given relating them to the Kauffman bracket skein module of a 3-manifold. The first theorem covers the case when the 3-manifold is split along a separating closed orientable surface and the second theorem addresses the case when the surface is nonseparating.

Keywords: skein module, three-manifold

1. Introduction

The study of knots and links was invigorated in 1984 by the introduction of the Jones polynomial [1]. It allowed solution of many long standing problems in knot theory including the Tate conjectures. One drawback was that the Jones polynomial is hard to compute for complicated knots and links. Then in 1987 Kauffman introduced the Kauffman bracket polynomial [2]. Easier to compute than the Jones polynomial, it carries essentially the same information (modulo framing and orientation). The fact that the Kauffman bracket takes framings into account makes it an ideal tool for studying links in arbitrary 3-manifolds, which need not have a canonical framing.

A major question at this time was how to construct something similar to the Jones polynomial for links in any 3-manifold. One approach was via topological quantum field theory. Another approach was through skein modules, introduced by Przytycki in 1989. [3] The importance of skein module theory was demonstrated when it was unified with topological quantum field theory by Blanchet-Habiger-Masbaum-Vogel [4], Lickorish [5], and Kauffman-Lins [6].

In this paper we study the Kauffman bracket skein module $K(X)$ of an oriented 3-manifold X . $K(X)$ has been computed for few X . In a paper by Hoste and Przytycki a method for computing $K(X)$ based on a handle decomposition of X was introduced. They found $K(X)$ when X is a lens space [7] and when $X = S^2 \times S^1$ [8]. Bullock used this approach to determine $K(X)$ when X is the complement of a $(2, 2p + 1)$ -torus knot [9]. Unfortunately this method seems too complicated when X has high Heegaard genus.

Until now, one thing that has been missing from the theory of Kauffman bracket

skein modules is a decomposition theorem in the spirit of the Mayer-Vietoris theorem from homology theory. Herein this problem is addressed. Let M be a smooth oriented 3-manifold and F a subsurface of ∂M . Informally the n th relative Kauffman bracket skein module of M , $K_n(M)$, is the Kauffman bracket skein module of M where the links have n arc components ending in $2n$ prescribed arcs in F . Suppose we have two manifolds like M , say X_1 and X_2 , and $X = X_1 \cup_F X_2$. We define a homomorphism

$$\Phi : \bigoplus_{n=0}^{\infty} (K_n(X_1) \otimes K_n(X_2)) \rightarrow K(X)$$

Intuitively, the map Φ is obtained by glueing an element of $K_n(X_1)$ to one of $K_n(X_2)$ to get an element of $K(X)$. It is shown that Φ is onto and its kernel is described simply and topologically.

Suppose L is a framed link in X intersecting F transversely in $2n$ arcs. We can view L as $L_1 \otimes L_2$ where $L_i = L \cap X_i$, $i = 1, 2$. Suppose that near F L_i is vertical in $F \times I$. If σ is a $2n$ component framed braid over F we can define $\sigma \cdot L_i$ as the result of replacing $L_i \cap (F \times I)$ with σ . The isotopy that takes $L_1 \otimes L_2$ to $\sigma \cdot L_1 \otimes \sigma^{-1} \cdot L_2$ is called a braiding move. Suppose some component of L_i bounds a bigon with one boundary component in L_i and the other in F . The isotopy pulling that part of L_i to the other side is a bigon move. The kernel is the submodule generated by all expressions of the form $L - L'$ where L' is obtained from L by a sequence of braiding and bigon moves.

2. Definitions

Suppose that X is an orientable three-manifold and F is a two-sided orientable surface embedded in X . Let $\psi : F \rightarrow F$ be a diffeomorphism and $X = X_1 \cup_{\psi} X_2$, where X_i is also an orientable three-manifold for $i = 1, 2$. Of course, we could choose coordinates so that ψ is the identity; we assume this to be the case throughout. We will define the relative Kauffman bracket skein modules for X_i , $K_n(X_i)$, and explore the relationship between the $K_n(X_i)$ and the Kauffman bracket skein module of X , $K(X)$.

Let M be an orientable three manifold with boundary $\partial M = F$. Let $a_1, \dots, a_{2n}$ be a collection of disjoint embedded arcs on F .

Definition 1. A framed arc in M is a homogeneous embedding of the square $I \times I$ into M with $(I \times I) \cap \partial M = (I \times \{0\}) \cup (I \times \{1\})$.

We call the parts of the framed arc lying on ∂M the ends of the arc, and denote it by $\partial_0 L$.

Usually when we draw framed arcs and circles we just draw them as arcs. In order to reflect the framing of the components, we use nugatory crossings and we take the framing to be the blackboard framing. See Figure 1 and Figure 2.

Definition 2. A collection of n framed arcs in M is nicely embedded if the ends of the framed arcs coincide with the a_j in a one-to-one manner.

We are now ready to define the relative Kauffman bracket skein modules of the X_i , but let's first recall the definition of the Kauffman bracket skein module of X .




Fig. 1. Two framed arcs




Fig. 2. Figure 1 represented as two arcs

Let R be the ring of Laurent polynomials over the integers and let $\mathcal{L}(X)$ be the set of all isotopy classes of links in X . The Kauffman bracket skein module of X is $R\mathcal{L}(X)/K$, where K is the submodule generated by the standard skein and frame relations. See Figure 3.

For the relative skein module we make the analogous definitions. That is, $\mathcal{L}_n(M)$ is the set of isotopy classes of nicely embedded framed links with n framed arcs and K_n is the submodule generated by the frame and skein relations with the relations also applied to the framed arcs. There is no generality lost in restricting to nicely embedded framed arcs in $\mathcal{L}_n(M)$, for there is an isotopy of any framed arc taking it to a nicely embedded one. A proof of this would be very much like the proof of Lemma 1.

Definition 3. The n^{th} relative Kauffman bracket skein module of M , written $K_n(M)$ is $R\mathcal{L}_n(M)/K_n$.

Let $B_{2n}(F)$ be the $2n$ -strand framed braid group over F . We want to define a group action of $B_{2n}(F)$ on $\mathcal{L}_n(M)$. Let $L \in \mathcal{L}_n(M)$. By taking a small product

$$\begin{array}{c}
\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = A \left(\begin{array}{c} \cup \\ \cap \end{array} \right) + A^{-1} \left(\begin{array}{c} \cap \\ \cup \end{array} \right) \\
\\
\begin{array}{c} \cup \\ \cap \end{array} = -A^{-3} \left(\begin{array}{c} \cap \\ \cup \end{array} \right)
\end{array}$$

Fig. 3. The skein and framing relations

neighborhood $\partial M \times I$ of ∂M in M it is possible to isotope L rel ∂M so that $L \cap (\partial M \times I) = (a_1 \times I) \cup \cdots \cup (a_{2n} \times I)$. Thus there is no harm in assuming this about L from the start.

Definition 4. Let $\sigma \in B_{2n}(F)$. Define $\sigma \cdot L = (L - (L \cap (\partial M \times I))) \cup \sigma$

Fig. 4. An example of the action of the $B_{2n}(F)$ on $\mathcal{L}_n(X_i)$

It is not hard to see that this really is a group action. It is clear that $(\sigma\tau) \cdot L = \sigma \cdot (\tau \cdot L)$, and $e \cdot L = L$ by the assumption on L immediately preceding the definition. Here e is the identity braid.

Let $L \in \mathcal{L}_n(M)$ and suppose that $H : (L \times I, \partial_0 L \times I) \rightarrow (M, \partial M)$ is an isotopy with $H_0(\partial_0 L) = H_1(\partial_0 L) = \{a_0, \dots, a_n\}$. This type of motion will be called a braiding move. Lemma 2 will allow us to see that $H_1(L) = \sigma \cdot L$ up to isotopy rel ∂M for some $\sigma \in B_{2n}(F)$, where L and $H_1(L)$ are considered as elements of $K_n(M)$.

Now suppose that L is a framed link in X and H is an isotopy of L in X with $H_0(L) \cap F = H_1(L) \cap F$, and $H_t(\partial_0 L) = H_t(L) \cap F$. Under these conditions $H_j(L) \cap X_i$ represents an element of $K_n(X_i)$ for $j = 0, 1$. Furthermore $H_1(L) \cap X_1 = \sigma \cdot (H_0(L) \cap X_1)$ and $H_1(L) \cap X_2 = \tau \cdot (H_0(L) \cap X_2)$ for some $\sigma, \tau \in B_{2n}(F)$. By choosing the right coordinates on ∂X_2 we would actually have $\tau = \sigma^{-1}$. Now let U_n be the submodule of $K_n(X_1) \otimes K_n(X_2)$ generated by all elements of the form

$$(a \otimes b) - (\sigma \cdot a \otimes \sigma^{-1} b)$$

Finally set $U = \bigoplus_{n=0}^{\infty} U_n$.

Definition 5. Suppose that $L \in \mathcal{L}_n(X_i)$ has an arc component c and there is a bigon bounded by c and some arc embedded in F . If L does not intersect the interior of this bigon then an isotopy that pulls c across to the other side of X and fixes the other components of L is called a bigon move.




Fig. 5. An example of a bigon move

Let V be the submodule of $\oplus_{n=0}^{\infty} [K_n(X_1) \otimes K_n(X_2)]$ generated by all elements of the form

$$(a \otimes b) - (c \otimes d)$$

where $c \otimes d$ is the result of applying a bigon move to $a \otimes b$.

Theorem 1. Let X be an orientable three-manifold, $F \subset M$ a closed orientable surface, and $\psi : F \rightarrow F$ a diffeomorphism. If $X = X_1 \cup_{\psi} X_2$ then

$$K(X) \cong \frac{\oplus_{n=0}^{\infty} K_n(X_1) \otimes K_n(X_2)}{U \vee V}$$

3. Lemmas

The first lemma we prove assures us that given any $L \in \mathcal{L}(X)$ it is represented by a link that intersects F in the prescribed way. Then we show that it is possible to reparameterize L so that, in the absence of bigon moves, an isotopy of L in X "splits in a nice way" into isotopies in X_1 and X_2 . Lastly we prove the theorem.

Lemma 1. Let $L \in \mathcal{L}_n(X)$. There is an isotopy of L so that $L \cap X_i \in \mathcal{L}_n(X_i)$, $i = 1, 2$ for some n .

Proof. Generically L intersects F in $2n$ arcs for some n . Let $s_1, \dots, s_{2n}$ be these arcs. Let t_i be one point of ∂a_i in F . By the Isotopy Lemma there is a diffeomorphism of F taking s_i to t_i for all i and this diffeomorphism is isotopic to the identity map on F . Call this isotopy $H : F \times I \rightarrow F$. Next let b be a smooth bump function on $[-1, 1]$ such that $b(-1) = b(1) = 0$ and $b(0) = 1$. We define another isotopy, called $\hat{H}$ on a small product neighborhood $F \times [-1, 1]$ of F in X . Let $\hat{H} : (F \times [-1, 1]) \times I \rightarrow [-1, 1]$ be given by $\hat{H}(x, s, t) = H(x, b(s)t)$. Finally it is clear that after applying $\hat{H}$ to L we can twist its band so that the altered L intersects F in the a_j . $\square$

Now let's focus our attention on the characterization of the submodules U_n .

Lemma 2. Let M be an orientable three-manifold with boundary F . Let $L \subset M$ be a properly embedded (finite) collection of disjoint framed arcs and framed circles. Suppose there are $n > 0$ framed arcs in L and $\partial L = \{a_1, \dots, a_{2n}\}$, where the a_j are as before. Suppose further that $H : L \times [0, 1] \rightarrow M$ is an isotopy




Fig. 6. The situation for the proof of Lemma 2

of L in M for which $H(L, 1) \cap \partial M = \{a_1, \dots, a_{2n}\}$. Then for some $\sigma \in B_{2n}(F)$, $H(L, 1) = \sigma \cdot L$ up to isotopy rel ∂M .

Proof. The strategy is to define a new isotopy $G : L \times [\delta, \eta] \rightarrow M$ equivalent to H in the sense that $G(L, \eta) = H(L, \eta)$. G will also have the property that $G(L, \frac{\delta+\eta}{2}) = \sigma \cdot L$ for $\sigma \in B_{2n}(F)$ and $G : L \times [\frac{\delta+\eta}{2}, 1] \rightarrow M$ is an isotopy rel ∂M . As in Lemma 1, we prove it for unframed nicely embedded links, but the result carries through.

We want to choose a product neighborhood $\partial M \times [-1, 1]$ with $\partial M = \partial M \times \{1\}$, so small that the arcs $L \cap (\partial M \times [-1, 1])$ have no critical points with respect to height. See Figure 6. Stated more precisely, let $f_s : [-\epsilon, \epsilon] \rightarrow \partial M \times [-1, 1]$ be a parametrization of a component of $(\partial M \times [-1, 1])$ at time s with $f_s(-\epsilon) \in \partial M$. Let $h : \partial M \times [-1, 1] \rightarrow [0, 2]$ be given by $h(x, t) = 1 - t$. Then we choose a neighborhood so small that $h \circ f_s$ is strictly increasing for all s .

Our goal is to show that we lose no generality in assuming that H is level-preserving in $\partial M \times [0, 1]$. This happens in two steps. First we make it preserve the $\partial M \times \{0\}$ level and then the $\partial M \times (0, 1)$ levels. We do this by reparametrizing the arcs $L \cap (\partial M \times [0, 1])$ one at a time. Denote by t the elements of $[-\epsilon, \epsilon]$ and by s those of $[\delta, \eta]$

For the first step define a function $\tau : [\delta, \eta] \rightarrow [-\epsilon, \epsilon]$ by

$$\tau(s) = f_s^{-1}(f_s([-\epsilon, \epsilon]) \cap (\partial M \times \{0\}))$$

This is well-defined since $h \circ f_s$ is increasing. We want a function $F : [-\epsilon, \epsilon] \times [\delta, \eta] \rightarrow [-\epsilon, \epsilon]$ so that

1. F_s is a diffeomorphism for all s , and
2. $F_s(\tau(s)) = 0$

Choose a small $\gamma > 0$ so that $(\tau(s) - \gamma, \tau(s) + \gamma) \subset [-\epsilon, \epsilon]$ for each s . Let $\beta_s : [-\epsilon, \epsilon] \rightarrow [0, 1]$ be a smooth function with $\beta_s(\tau(s)) = 1$ and $\beta_s(t) = 0$ outside of

$(\tau(s) - \gamma, \tau(s) + \gamma)$. Then define $F_s(t) = t - \tau(s)\beta_s(t)$. Clearly $F_s(\tau(s)) = 0$. The proof that F_s is a diffeomorphism is rather technical and is omitted. The desired parametrizations are given by $g_s = f_s \circ F_s^{-1}$.

Now we want to make H preserve levels in $\partial M \times [0, 1]$. To this end we reparametrize again so that each arc is parametrized by height. Fortunately this is much easier than the first part. Notice that for $h \circ g_s : [-\epsilon, 0] \rightarrow [0, 1]$ we have

1. $h \circ g_s(-\epsilon) = 0$
2. $h \circ g_s(0) = 1$, and
3. $(h \circ g_s)'(t) \neq 0$ for all t

Thus $h \circ g_s$ is a diffeomorphism for each s . Let $\zeta_s : [0, 1] \rightarrow [-\epsilon, 0]$ be $(h \circ g_s)^{-1}$, and consider $g_s \circ \zeta_s : [0, 1] \rightarrow M$, a parametrization of one of the arcs. Since $h \circ g_s \circ \zeta_s = 1_{[0, 1]}$ the arc is parametrized by height for all s . Clearly we can do this for each arc in $\partial M \times [0, 1]$, so there is no harm in assuming that H is level preserving in a neighborhood of ∂M .

Next we define G . For ease of notation let's assume that $\delta = 0$ and $\eta = 1$. Let $b : I \rightarrow I$ be a smooth function with $b(0) = 1$ and $b(1) = 0$. Then make

$$G(x, t) = \begin{cases} x & x \in M - (\partial M \times I), 0 \leq t \leq \frac{1}{2} \\ H(x, 2b_1(s)t) & x = (y, s) \in \partial M \times I, 0 \leq t \leq \frac{1}{2} \\ H(x, 2t - 1) & x \in M - (\partial M \times I), \frac{1}{2} < t \leq 1 \\ H(x, (2 - 2b_1(s))(t - 1) + 1) & x = (y, s) \in \partial M \times I, \frac{1}{2} < t \leq 1 \end{cases}$$

It is easy to check that $G(L, 1) = H(L, 1)$ and that the maps line up properly at $t = \frac{1}{2}$ and at the boundary of $\partial M \times I$ that is interior to M . Notice that the only way that G differs from H is in the speed at which the isotopy happens. This means that our assumption that H is level preserving assures us that $H(L, t)$ is actually an embedding for all $t \in I$. Also it isn't hard to see that $G(L, \frac{1}{2}) = \sigma \cdot L$ and G fixes ∂M for all $t \in [\frac{1}{2}, 1]$. $\square$

Ying-Qing Wu suggested a much shorter proof of Lemma 2 but in the proof of the main theorem we want to use this technique again, so we use the longer version.

4. Proof of the Theorem

Proof. The first step is to define an R -module homomorphism from $\mathcal{L}_n(X_1) \times \mathcal{L}_n(X_2)$ to $\mathcal{L}(X)$. If we just take the union the framed arcs become framed circles. Call this map q_n . It is well-defined since the framed braids on either side are nicely embedded. Extend this to a map

$$q_n : R\mathcal{L}(X_1) \times R\mathcal{L}(X_2) \rightarrow R\mathcal{L}(X)$$

bilinearly. This induces a homomorphism

$$\hat{q}_n : R\mathcal{L}(X_1) \otimes R\mathcal{L}(X_2) \rightarrow R\mathcal{L}(X)$$

Now let S be the submodule of $R\mathcal{L}(X_1) \otimes R\mathcal{L}(X_2)$ generated by all elements of the form $a' \otimes b$ or $a \otimes b'$, where $a \in R\mathcal{L}(X_1)$, $b \in R\mathcal{L}(X_2)$, $a' \in K_n^1$, and $b' \in K_n^2$.

(Recall that K_n^i is the smallest submodule of $R\mathcal{L}_n(X_i)$ containing all of the skein and framing relations.) Notice that $\hat{q}_n(S) \subset K$. Therefore there is a homomorphism

$$\tilde{q}_n : [R\mathcal{L}(X_1) \otimes R\mathcal{L}(X_2)]/S \rightarrow K(X)$$

But $(R\mathcal{L}(X_1) \otimes R\mathcal{L}(X_2))/S \cong K_n(X_1) \otimes K_n(X_2)$. So we get a map

$$\tilde{q}_n : K_n(X_1) \otimes K_n(X_2) \rightarrow K(X)$$

Thus for each n there are embeddings

$$\varphi_n : [K_n(X_1) \otimes K_n(X_2)]/\ker \tilde{q}_n \rightarrow K(X)$$

This implies the existence of

$$\Phi : \bigoplus_{n=0}^{\infty} [K_n(X_1) \otimes K_n(X_2)]/\ker \tilde{q}_n \rightarrow K(X)$$

By Lemma 1 Φ is onto

$$\begin{aligned} &\Rightarrow \bigoplus_{n=0}^{\infty} \frac{[K_n(X_1) \otimes K_n(X_2)]}{\ker \tilde{q}_n} / \ker \Phi \cong K(X) \\ &\Rightarrow \frac{\bigoplus_{n=0}^{\infty} [K_n(X_1) \otimes K_n(X_2)]}{\bigoplus_{n=0}^{\infty} \ker \tilde{q}_n} / \ker \Phi \cong K(X) \end{aligned}$$

This is a start but the result will not be very useful until we characterize $\bigoplus_{n=0}^{\infty} \ker \tilde{q}_n$ and $\ker \Phi$ topologically. The first will be described with the help of the group action discussed earlier and we will deal with the second via bigon moves.

We want to show that if $L \in \mathcal{L}(X)$ and $H : L \times I \rightarrow X$ is an isotopy then $H_0(L)$ and $H_1(L)$ are related by a sequence of braiding moves and bigon moves. There is no harm in assuming that $H_0(L) \cap F = \{a_1, \dots, a_{2n}\} = H_1(L) \cap F$. Furthermore we reparameterize the arcs near their intersection with F using the technique of the proof of Lemma 2. This allows us to view H as two separate isotopies; one on X_1 and one on X_2 . We also assume that the motion H is generic. This means that there exist $s_1, \dots, s_k \in I$ for which $|H_{s_i}(L) \cap F|$ is odd, and further only one of these points is non-generic.

Choose ϵ so small that $H : L \times [s_i - \epsilon, s_i + \epsilon] \rightarrow X$ is a bigon move. Of course this isn't technically the case since the other components of L move some, but by choosing ϵ small we can ignore this. The strategy is to speed up H so that what occurs from $s = 0$ to $s = s_0 - \epsilon$ occurs from $s = 0$ to $s = s_0 - 2\epsilon$. Then we "undo" H near F from $s = s_0 - 2\epsilon$ to $s = s_0 - \epsilon$. At this point we apply Lemma 3.2, which gives a braiding move. Then from $s = s_0 - \epsilon$ to $s = s_0 + \epsilon$ we have a bigon move. Then between $s = s_0 + \epsilon$ and $s = s_0 + 2\epsilon$ we put the link back to how it was at $s = s_0 - \epsilon$. Repeat this process for all of the s_i and we're done.

Let $u = \frac{s_0 - \epsilon}{s_0 - 2\epsilon}$ and define $G : L \times [0, s_0 - 2\epsilon] \rightarrow X$ by $F(x, s) = H(x, us)$. Then choose a small product neighborhood $F \times [-1, 1]$ of F in X with $F = F \times \{0\}$. Let $g : F \times I \rightarrow F$ be an isotopy of F taking $H_{s_0 - \epsilon}(L) \cap F$ to $H_0(L) \cap F$. Next




Fig. 7. The isotopy H

let $b : [-1, 1] \rightarrow [0, 1]$ be a smooth function with $b(0) = 1$ and $b(1) = b(-1) = 0$. Consider $G : (F \times [-1, 1]) \times [s_0 - 2\epsilon, s_0 + \epsilon] \rightarrow X$ given by

$$G(x, t, s) = (g(x, b(t)s), t)$$

. Then G satisfies the hypotheses of Lemma 3.2 and so is a braiding move. Extend G to $[0, s_0 + \epsilon]$ by tacking on the appropriate bigon move. Further extend G by taking

$$G(x, t, s) = (g(x, 1 - b(t)s), t)$$

for $s \in [s_0 + \epsilon, s_0 + 2\epsilon]$. Finally repeat this process as many times as necessary to see that $H_1(L)$ as $H_0(L)$ with a sequence of braiding moves and bigon moves applied to it. $\square$

5. The Case When F is Nonseparating

We now assume tha X and F are as before, but F is nonseparating. In other words, $X = X_1/\psi$ where $\psi : F \rightarrow F$ is the diffeomorphism and X_1 is X cut along F . With some minor changes in the definitions the theorem carries over to this setting. What's more, the same proof works. After cutting X along F , X_1 contains two copies of F , say $F, F' \subset \partial X_1$. This gives us two sets of embedded arcs, $a_j \subset F$, and $a'_j \subset F'$.

Definition 6. A collection of framed arcs in X_1 is nicely embedded if the ends of the framed arcs coincide with the a_j and the a'_j in a one- to-one manner.

Now we discuss the necessary changes in the definition of braiding moves. Here we define both a left and a right action of $B_n(F)$ on $\mathcal{L}_n(X_1)$. The left action is the same as in definition 4 and is again written $\sigma \cdot L$. The right action is exactly the same but it happens near F' and is written $L \cdot \sigma$. (Here $L \in \mathcal{L}_n(X_1)$ and $\sigma \in B_n(F) \cong B_n(F')$.) For each n let U_n be the submodule of $K_n(X)$ generated by all the elements of the form $a - \sigma \cdot a \cdot \sigma^{-1}$, where $a \in K_n(X_1)$ and $\sigma \in B_n(F)$. Then let $U = \oplus_{n=0}^{\infty} U_n$ and call this the submodule generated by all braiding moves.

The definition of a bigon move is the same as before. Here we define V to be the submodule of $K_n(X_1)$ generated by all elements of the form $a - a'$ where a' is the result of applying a bigon move to a .

Theorem 2. Let X be an orientable 3-manifold, $F \subset X$ a closed, orientable nonseparating surface, and $\psi : F \rightarrow F$ a diffeomorphism. Let X_1 be X cut along F , so that $X = X_1/\psi$. Then

$$K(X) \cong \frac{\oplus_{n=0}^{\infty} K_n(X_1)}{U \vee V}$$

Proof. This proof differs from that of Theorem 1 only in the algebraic part. Obviously there is no tensor product here. Nonetheless this part of the proof is the same in spirit. In lieu of writing it out, we just say that the map Φ is defined from $\oplus_{n=0}^{\infty} K_n(X_1)$ instead of from $\oplus_{n=0}^{\infty} K_n(X_1) \otimes K_n(X_2)$. The rest of the proof is identical since it all occurs near the surface F . $\square$

References

- [1] V.F.R Jones *A polynomial invariant for knots via Von Neuman algebras*, Bull. AMS. **12** (1985) 103-11.
- [2] L. Kauffman *State models and the Jones polynomial*, Topology **26** no. 3 (1987) 395-401
- [3] J.H. Przytycki, *Skein modules of 3-manifolds*, Bull. Pol. Acad. Sci. **39(1-2)** (1991) 91-100.
- [4] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel *Topological quantum field theories derived from the Kauffman bracket*, Topology, **34** no.4 (1995) 883-927
- [5] W.B.R. Lickorish *The skein method for three-manifold invariants*, J. Knot Theory Ramifications **2** no. 2 (1993) 171-194
- [6] L. Kauffman and S.L. Lins, *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, Annals of Math. Studies, Princeton University Press (1994)
- [7] J. Hoste and J.H. Przytycki, *The $(2, \infty)$ -skein module of lens spaces; a generalization of the jones polynomial*, J. Knot Theory Ramifications **2** no. 3 (1993) 321-333.
- [8] J. Hoste and J.H. Przytycki, *The Kauffman bracket skein module of $S^2 \times S^1$* , Math Z. **220** (1995) 65-73.
- [9] D. Bullock, *The $(2, \infty)$ -skein module of the complement of a $(2, 2p+1)$ torus knot*, J. Knot Theory Ramifications **4** no. 4 (1995) 619-632.